

IMMUNIZATIONS, VACCINATIONS, AND OTHER MISCELLANEOUS TREATMENTS	DEFER FOR:
ABRYSVO (RSV vaccine)	Accept immediately if feeling well and healthy
Allergy Desensitization	Accept immediately if feeling well and healthy
Anthrax	Accept immediately if feeling well and healthy
AREXVY (RSV vaccine)	Accept immediately if feeling well and healthy
Botulinum Toxin ('Botox')	Defer for 24 hours after treatment
BCG	2 weeks
Chicken Pox (Varivax)	4 weeks
Chicken Pox (VZIG) Immune Globulin	12 months
Cholera	Accept immediately if feeling well and healthy
COVID-19	<ul style="list-style-type: none"> • Accept immediately if feeling well and healthy after receiving a Non-Replicating, Inactivated, Viral Vector (such as Johnson & Johnson), mRNA (such as Pfizer/BioNTech, Comirnaty or Moderna) or subunit protein (such as Novavax) vaccine that has been approved in the USA • 2 weeks if feeling well and healthy after receiving a vaccine that has not been approved and is in clinical trials in USA, Europe, Canada, or Australia (such as AstraZeneca (Covishield)/Oxford, CureVac, and Sanofi/GSK) • 2 weeks If the donor does not remember the name of the vaccine and received it in USA, Europe, Canada, or Australia • 1 year if feeling well and healthy after receiving a vaccine in use or in clinical trials in countries other than USA, Europe, Canada, or Australia (such as Cansino, Bharat, Gamaleya, Sinopharm (Beijing), Sinopharm (Wuhan), Sinovac Biotech, Sputnik V and Vector Institute) • 1 year if the donor does not remember the name of the vaccine and received it in a country other than USA, Europe, Canada, or Australia • Refer to NY-JA-0521 "COVID-19 Vaccine List with Deferral Intervals" for COVID-19 vaccinations not listed
Diphtheria	Accept immediately if feeling well and healthy
DT: Diphtheria \ Tetanus	Accept immediately if feeling well and healthy

IMMUNIZATIONS, VACCINATIONS, AND OTHER MISCELLANEOUS TREATMENTS	DEFER FOR:
DPT: Diphtheria \ Pertussis \ Tetanus	Accept immediately if feeling well and healthy
Flu (injection or nasal)	Accept immediately if feeling well and healthy
Gamma Globulin - prophylactic	Accept immediately if feeling well and healthy and given for general prophylaxis (e.g., for travel) and not for exposure
Gamma Globulin - exposure	<ul style="list-style-type: none"> • 120 days (4 months) if given as a precaution to Hepatitis A exposure (i.e., restaurant outbreak) • Evaluate if given for exposure to any other disease
Hepatitis A - prophylactic	Accept immediately if feeling well and healthy
Hepatitis A - exposure	120 days (4 months)
Hepatitis B - prophylactic	2 weeks
Hepatitis B - exposure	3 months
Hep A + Hep B - prophylactic	2 weeks
Hep A + Hep B - exposure	4 months
HIB: Haemophilus Influenzae B	Accept immediately if feeling well and healthy
HIV Prevention (PrEP and PEP): Apretude \ cabotegravir	2 Years following date of last injection or shot
Human Papilloma Virus	Accept immediately if feeling well and healthy
IVIG (Intravenous Immune Globulin)	Defer for 12 months
Japanese Encephalitis	Accept immediately if feeling well and healthy
Lyme	Accept immediately if feeling well and healthy
Measles (Rubeola)	2 weeks
Measles, German (Rubella)	4 weeks
Meningococcus (including Meningitis B)	Accept immediately if feeling well and healthy
MMR (Measles, Mumps, Rubella)	4 weeks
Mpox (Monkeypox)	<ul style="list-style-type: none"> • Jynneos prophylactically: Accept immediately if feeling well and healthy • Jynneos post exposure, no complication: 21 days • Jynneos post exposure, with complication: 1 month or upon resolve of symptoms, whichever is later • ACAM2000 (not FDA licensed): 8 weeks
Mounjaro (tirzepatide) injection	Accept immediately if feeling well and healthy

IMMUNIZATIONS, VACCINATIONS, AND OTHER MISCELLANEOUS TREATMENTS	DEFER FOR:
Mumps	2 weeks
Ozempic (semaglutide) injection	Accept immediately if feeling well and healthy
Paratyphoid	Accept immediately if feeling well and healthy
Pertussis	Accept immediately if feeling well and healthy
Plague	Accept immediately if feeling well and healthy
Pneumonia	Accept immediately if feeling well and healthy
Polio: Oral	2 weeks
Polio: Injection	Accept immediately if feeling well and healthy
Rabies	<ul style="list-style-type: none"> • Accept immediately if received for prophylaxis (e.g., animal care worker) and not for exposure • Defer 12 months if due to animal bite
Rocky Mountain Spotted Fever	Accept immediately if feeling well and healthy
Rybelsus (semaglutide) tablets	Accept immediately if feeling well and healthy
Shingles: Zostavax	4 weeks
Shingles: Shingrix (Zoster Vaccine Recombinant)	Accept immediately if feeling well and healthy
Smallpox without complications	8 weeks
Smallpox with complications	<ul style="list-style-type: none"> • 8 weeks after onset of symptoms or 14 days after symptoms resolve, whichever is longer • Complications include the following: Rash (resembling ‘blisters’) covering a small or large area of the body; necrosis (tissue death) in the area of vaccination; inflammation of the brain (encephalitis); infection of the cornea (eye); localized or systemic skin reaction in someone with eczema or other chronic skin condition
Tetanus Toxoid (Booster)	Accept immediately if feeling well and healthy
Tetanus and Diphtheria Toxoids	Accept immediately if feeling well and healthy
Typhoid: Injection	Accept immediately if feeling well and healthy
Typhoid: Oral	2 weeks
Typhus, Paratyphoid	Accept immediately if feeling well and healthy
Wegovy (semaglutide) injection	Accept immediately if feeling well and healthy
Yellow Fever	2 weeks